

Tongue depressor as an intubation aid in a patient with vallecular cyst: A case report

Abdul Monem¹, Fauzia Anis Khan², Sohail Awan³

Abstract

Introduction: Vallecular cyst is a rare occurring often encountered incidentally, which may compromise the airway during induction of anesthesia. There is a high risk for cyst rupture and soiling of the airway during manipulation of the airway.

Case Report: We report a case of a young 23-year-old female with a history of difficulty in breathing and swallowing for the past 1½ years. She underwent emergency tracheostomy and excision of some oral swelling 10 years back. She was electively booked for excision of the cyst. Various techniques and instruments have been described in the past. We used a tongue depressor to gently push the cyst aside and successfully rail loaded a reinforced endotracheal tube over a gum elastic bougie.

Conclusion: Intubation in a patient with oropharyngeal cyst poses a significant risk in view of complete airway obstruction or aspiration of the contents in case rupture of the cyst. Various techniques for intubation have been reported, either awake or under general anesthesia. We successfully used a tongue depressor to shift the cyst away as an aid for intubation. This technique has so far not been reported in the literature.

Keywords: Vallecular cyst, induction, tongue depressor.

Background

Vallecular cyst is a rare cause of airway collapse, which may remain unidentified till induction of general anesthesia. Various methods have been reported to handle such cases. We used a tongue depressor and a gum elastic bougie as an aid to intubation in a 23-year-old female.

Case Report

A 23-year-old female presented with a history of snoring and occasional episodes of dysphagia and difficulty in breathing for the past 1½ years. There was no history of change in symptoms associated with a change in body position. She also had a history of palpitations for which she was prescribed propranolol by a general physician. Her thyroid function tests were normal. She had been operated twice before

at the age of 10 and 13 years, for a neck swelling which had caused difficulty in breathing and swallowing. She underwent tracheostomy during the first surgery. At the ear, nose, and throat clinic, she underwent an indirect laryngoscopy. A 3 cm × 3 cm swelling was seen arising from the right side of the vallecula. Vocal cords were visible after the fiber-optic scope was negotiated below the swelling. Both cords were normal in anatomy and equally mobile. Her pre-operative computed tomography scan is given in Fig. 1. She was labeled the American Society of Anesthesiologist II with normal laboratory findings including normal thyroid function tests. She had a Grade I Mallampati score.

Perioperative course

She was planned for excision of cyst under

general anesthesia with consent for tracheostomy if required. She underwent inhalational induction with sevoflurane 1–8% in 100% oxygen. Careful laryngoscopy was performed using a size 3 Macintosh blade after attaining adequate depth of anesthesia. After the ability to hand ventilate was ensured, intravenous bolus of 100 µg fentanyl and 50 mg of propofol was administered. Laryngoscopy revealed a large cyst obscuring the glottic opening (Fig. 2). A metallic tongue depressor was used to gently press and shift the cyst to the right. A narrow slit-like glottic opening then became visible. A size 6 reinforced tracheal tube was then negotiated over a well-lubricated gum elastic bougie without using a muscle relaxant (Fig. 3). Tracheal tube was fixed at 20 cm mark after confirming bilateral air entry and end-tidal capnography. There was no episode of desaturation during the procedure. Muscle relaxation was provided using atracurium 0.5 mg/kg. After inserting a throat pack, the surgeon excised the 3 cm × 3 cm cyst along with the right tonsil (Fig. 4). Two similar cysts were also excised, one through right lateral pharyngotomy and another from the front of the neck (Fig.

¹Department of Anaesthesiology, Aga Khan University Hospital

²Department of Anaesthesiology, Aga Khan University Hospital

³Department of Surgery, Aga Khan University Hospital

Address of Correspondence

Dr. Dr. Abdul Monem,

Department of Anaesthesiology, Aga Khan University

E-mail: abdul.monem@aku.edu



Dr. Abdul Monem



Dr. Fauzia Anis Khan



Dr. Sohail Awan

© 2018 by Journal of Anaesthesia and Critical Care Case Reports | Available on www.jaccr.com |

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.




Figure 1: Pre-operative computed tomography view showing the vallecular cyst.




Figure 2: Laryngoscopic view with the cyst **Figure 3:** After intubation. **Figure 4:** After excision.

5). Patient's trachea was extubated after insertion of a nasogastric tube. Her post-operative recovery was uneventful and she was discharged 2 days later. Biopsy reported the specimen to be a lipoma.

Discussion

Congenital laryngeal cysts are rare, with an incidence of 1.82 per 100,000 live births [1]. Common sites are 58.2% in the glottic area and 18.3% in the ventricular folds. The remainder was located on the aryepiglottic fold (2.2%) and interarytenoid region (0.7%) [2]. These are benign and do not usually cause problems, but unidentified cases are very prone to sudden severe hypoxia and death due to complete airway collapse, especially in children. Adults may present with symptoms of change in voice or dysphagia. Laryngocele and saccular cysts are also rare cause of such symptoms. Laryngoceles cause episodic symptoms as they intermittently fill with air, whereas saccular cysts cause constant symptoms. Children usually present with difficulty in breathing, swallowing, and failure to thrive. Yadav [3] reported hemoptysis as a presenting symptom. Trauma and cyst rupture may lead to aspiration during intubation. Mason and Wark [4] reported two cases of difficult mask ventilation after intravenous induction. Both were found to have vallecular cyst obscuring the glottis view on laryngoscopy. One was intubated after successful aspiration of the cyst by the ENT surgeon. Other was intubated with the help of a bougie. McHugh [5] also reported

successful intubation with the help of a bougie in an unexpected case of a large epiglottic cyst. Different methods of airway maintenance have been reported in the anesthetic management of these cases. McKiernan [6] reported blind intubation in a 7-year-old child with the help of smaller tube with double bend on a malleable introducer by pushing an unexpected vallecular cyst to one side. Awake fiber-optic intubation is considered the safest technique in difficult airway management, but this technique may be associated with difficulty to negotiate around the cyst. It may also cause rupture of cyst or traumatic bleed leading to aspiration. In spite of these disadvantages, it has been successfully used in infants [7]. Ahrenset al. [8] reported intubation of a 3-month-old child with a 2.2 fiber-optic bronchoscope through a laryngeal mask airway. Remifentanyl and propofol sedation has been used for fiber-optic intubation in an obstetric patient with a vallecular cyst [9]. It does not increase the risk of aspiration in the mother or leads to sedation in the newborn [10]. Our patient was too anxious to undergo awake fiber-optic intubation. We did not want to administer sedatives in our patient for fear of airway obstruction, as she was able to maintain the airway under inhalational anesthesia, and use of fiberoptic laryngoscope was an alternate method that could have been used after inhalational induction. This was our plan B if the use of tongue depressor had failed. Other methods to negotiate the endotracheal tube that has been used include shifting the cyst to one side with the help of a styletted endotracheal tube [11] or lifting of the cyst with a larger Macintosh or a rigid bronchoscopy blade. An angled atraumatic instrument like Mac gills forceps could also be used to shift the cyst away from the intubation path. Although in this case the cyst was thick walled and there was less likelihood of rupture, in case of thin cysts,

two wide bore suctions ought to be ready. Shifting the patient to lateral position might help to reopen the complete airway collapse at induction, especially in children [12]. The use of backward, upward, rightward pressure maneuver has been described in a 4-month-old infant [13]. Consent should also be taken for standby tracheostomy. Marsupialization of the vallecular cyst under local anesthesia has also been reported by Pagella et al. [14]. In our case, we used a tongue depressor to push the cyst so that a view of laryngeal inlet could be obtained. By doing so, we were able to intubate the trachea with a size 6 polyvinyl chloride in forced tracheal tube over a gum elastic bougie. There was some resistance while inserting the tube probably because no muscle relaxant had been used. The use of 4% lignocaine spray might have eased intubation in this patient. The use of tongue depressor in the management of vallecular cyst has not been reported in literature. This simple method was helpful in our case mainly because the cyst was firm and could be displaced laterally. If the cyst had been soft or pedunculated, this might not have been possible. If general anesthesia is chosen as induction technique, inhalational induction with careful laryngoscopic examination should precede attempt at intubation. As the use of muscle relaxants may cause complete airway obstruction, they should be administered after successful endotracheal intubation. Surgical help to perform tracheostomy or puncture of the cyst should be readily available for which consent should be taken before hand.

Conclusion

We recommend this simple method of using a tongue depressor as a tracheal intubation aid in anesthetic management of a patient with vallecular cyst. Other atraumatic intubation aids and external laryngeal maneuver might be helpful as well and a gum elastic bougie must be handy in such



Figure 5: Removed cysts.

situations. Consent for emergency tracheostomy should also be taken.

References

1. Pak MW, Woo JK, Van Hasselt CA. Congenital laryngeal cysts: Current approach to management. *J Laryngol Otol* 1996;10:854-6.
2. Arens C, Glanz H, Kleinsasser O. Clinical and morphological aspects of laryngeal cysts. *Eur Arch Otorhinolaryngol* 1997;254:430-6.
3. Yadav SP, Sahni JK, Raj B, Chawla RK, Yadav J, Singh B. Laryngeal cyst causing haemoptysis. An unusual presentation. *Indian J Chest Dis Allied Sci* 1986;28:50.1.
4. Mason DG, Wark KJ. Unexpected difficult intubation. Asymptomatic epiglottic cysts as a cause of upper airway obstruction during anaesthesia. *Anaesthesia* 1987;42:407-12.
5. McHugh P. Cyst of epiglottis (letter). *Anaesthesia* 1989;44:522.
6. McKiernan EP, Meakin G. Vallecular cysts (letter). *Anaesthesia* 1988;43:808-9.
7. Cheng KS, Ju-Mei Ng, Li HY, Hartigan PM. Vallecular cyst and laryngomalacia in infants: Report of six cases and airway management. *Anesth Analg* 2002;95:1248-50.
8. Ahrens B, Lammert I, Schmitt M, Wahn U, Paul K, Gemann B. Life-threatening vallecular cyst in a 3-month-old infant: Case report and literature review. *Clin Pediatr (Phila)* 2004;43:287-90.
9. Prabhu A, ManNally L, Pradhan P, Amin HH. Awake Fiberoptic Intubation in a Pregnant Patient with a Laryngeal Cyst. *Int J Periop Ultrasound Appl Technol* 2012;1:74-6.
10. Ovassapian A, Krejcie TC. Awake fiberoptic intubation in a patient at high risk of aspiration. *BJA* 1989;62:12.
11. Kamble VA, Lilly RB, Gross JB. Unanticipated difficult intubation as a result of an asymptomatic vallecular cyst. *Anaesthesiology* 1999;3:872-3.
12. Walshe CM, Jonas N, Rohan D. Vallecular cyst causing a difficult intubation. *Br J Anaesth* 2009;102:565.
13. Namshikar VN, Dukle NV, Sukhthanker DS. Anesthetic management of vallecular cyst excision in an infant: An airway challenge: Case report. *Saudi J Anesth* 2016;10:356-8.
14. Pagella F, Pusateri A, Matti E, Tinelli G, Benazzo M. Transoral power assisted Marsupialization of vallecular cyst under local anaesthesia. *Laryngoscope* 2013;123:699-701.

Conflict of Interest: Nil
Source of Support: None

How to Cite this Article

Monem A, Khan F A, Awan S. Tongue depressor as an intubation aid in a patient with vallecular cyst: A case report. *Journal of Anaesthesia and Critical Care Case Reports* Jan-Apr 2018; 4(1):16-18.